# DEPARTMENT of ENVIRONMENTAL SERVICES Water Supply & Pollution Control Division - Biology Bureau

## LAKE TROPHIC DATA

### MORPHOMETRIC:

Lake: BAKER POND		Lake Area (ha):	5.75
Town: CHESTERF	IELD	Maximum depth (m):	7.1
County: Cheshire		Mean depth (m):	2.2
River Basin: Connecti	cut	Volume (m³):	124500
Latitude: 42°51'08		Relative depth:	2.6
Longitude: 72°27'59		Shore configuration:	1.18
Elevation (ft):	1130	Areal water load (m/yr)	: 15.28
Shore length (m):		Flushing rate $(yr^{-1})$ :	7.10
Watershed area (ha):	173.5	P retention coeff.:	0.50
<pre>% watershed ponded:</pre>	0.0	Lake type:	natural

BIOLOGICAL:	9 March 1994	30 August 1993
DOM. PHYTOPLANKTON (% TOTAL) #1	DINOBRYON 35%	DINOBRYON 99%
#2	TINY PENNATE DIATOMS 308	
#3	(ALL ALGAE SPARSE)	
PHYTOPLANKTON ABUNDANCE (cells/mL)		570
CHLOROPHYLL-A (µg/L)		7.55
DOM. ZOOPLANKTON (% TOTAL) #1	SPARSE ZOOPLANKTON	NAUPLIUS LARVA 64%
#2	ONLY TWO DAPHNIA	
#3	OBSERVED	
ROTIFERS/LITER	<1	8
MICROCRUSTACEA/LITER	2	47
ZOOPLANKTON ABUNDANCE (#/L)	2	55
VASCULAR PLANT ABUNDANCE		Sparse
SECCHI DISK TRANSPARENCY (m)		3.5
BOTTOM DISSOLVED OXYGEN (mg/L)	6.0	0.4
BACTERIA (E. coli, #/100 ml) #1		
#2		
#3		

### SUMMER THERMAL STRATIFICATION:

# weakly stratified

Depth of thermocline (m): None Hypolimnion volume (m³): None

Anoxic volume  $(m^3)$ : 350

CHEMICAL:			BAKER PON CHESTERFI		
	9 March	n 1994	30 August 1993		93
DEPTH (m)	2.0	5.5	2.5		5.0
pH (units)	4.9	5.1	5.2		5.3
A.N.C. (Alkalinity)	0.2	0.7	0.0		0.1
NITRATE NITROGEN	0.08	0.07	< 0.02		< 0.02
TOTAL KJELDAHL NITROGEN	0.63	0.71	0.47		0.50
TOTAL PHOSPHORUS	0.002	0.006	0.017		0.011
CONDUCTIVITY (µmhos/cm)	28.4	27.9	23.1		23.5
APPARENT COLOR (cpu)	11	17	10	· · · · · · · · · · · · · · · · · · ·	12
MAGNESIUM			0.29		
CALCIUM			1.0		
SODIUM			1.2		
POTASSIUM			< 0.40		
CHLORIDE	< 2	< 2	< 3		< 3
SULFATE	7	7	6		6
TN : TP	355	130	28		45
CALCITE SATURATION INDEX					


All results in mg/L unless indicated otherwise

TROPHIC CLASSIFICATION: 1993

D.O.	S.D.	PLANT	CHL	TOTAL	CLASS
**	2	0	1	3	Oligo.

#### **COMMENTS:**

- 1. Also known as Baker-Hubbard Pond.
- 2. This is a remote pond located within Pisgah State Park, an undeveloped, wilderness park. It was sampled cooperatively with the NH Fish and Game Department.
- 3. This is a non-colored acid pond with very low buffering capacity (A.N.C.).
- 4. The dominant genera of wholewater phytoplankton were <u>Dinobryon</u> (60%), <u>Cryptomonas</u> (20%) and <u>Mallomonas</u> (15%).



#### FIELD DATA SHEET

LAKE: BAKER POND TOWN: CHESTERFIELD

DATE: 08/30/93 WEATHER: SUNNY, WARM


DEPTH (M)	TEMP (°C)	*DISSOLVED OXYGEN	OXYGEN SATURATION	
0.5	25.5	8.1	98 %	
1.0	25.0	8.0	97 %	
2.0	25.0	8.0	97 %	
3.0	24.5	7.7	90 %	
4.0	23.5	5.6	65 %	
5.0	23.0	3.8	44 %	
6.0	22.5	1.4	16 %	
6.5	22.5	0.4	5 %	

SECCHI DISK (m): 3.5 COMMENTS:

BOTTOM DEPTH (m): 7.1

TIME: 1145

\*Dissolved oxygen values are in mg/L



## AQUATIC PLANT SURVEY

LAK	E: BAKER POND	TOWN: CHESTERFIELD	DATE: 08/30/93	
Vor	PLANT	NAME		
Кеу	GENERIC	COMMON	ABUNDANCE	
N	Nymphaea	White water lily	Sparse	
υ	Utricularia	Bladderwort	Sparse	
P	Pontederia cordata	Pickerelweed	Sparse	
Y	Nuphar	Yellow water lily	Sparse	
d	Dulichium arundinaceum	Three-way sedge	Sparse	
х		Sterile thread-like leaf	Sparse	
-				
	•			
	- · · · · · · · · · · · · · · · · · · ·			
	4		<u> </u>	
		OVERALL ABUNDANCE:	Sname o	

# **GENERAL OBSERVATIONS:**

1. Pond was surrounded by leatherleaf, and leatherleaf was present on boggy islands. Because the leatherleaf was not actually in the water, it is not depicted on the map.